

Nutrition and Health – The Association between Eating Behavior and Various Health Parameters: A Matched Sample Study

Nathalie T. Burkert, Johanna Muckenhuber, Franziska Großschädl, Éva Rásky, Wolfgang Freidl

Published: February 7, 2014 • <https://doi.org/10.1371/journal.pone.0088278>

Abstract

Population-based studies have consistently shown that our diet has an influence on health. Therefore, the aim of our study was to analyze differences between different dietary habit groups in terms of health-related variables. The sample used for this cross-sectional study was taken from the Austrian Health Interview Survey AT-HIS 2006/07. In a first step, subjects were matched according to their age, sex, and socioeconomic status (SES). After matching, the total number of subjects included in the analysis was 1320 (N=330 for each form of diet – vegetarian, carnivorous diet rich in fruits and vegetables, carnivorous diet less rich in meat, and carnivorous diet rich in meat). Analyses of variance were conducted controlling for lifestyle factors in the following domains: health (self-assessed health, impairment, number of chronic conditions, vascular risk), health care (medical treatment, vaccinations, preventive check-ups), and quality of life. In addition, differences concerning the presence of 18 chronic conditions were analyzed by means of Chi-square tests. Overall, 76.4% of all subjects were female. 40.0% of the individuals were younger than 30 years, 35.4% between 30 and 49 years, and 24.0% older than 50 years. 30.3% of the subjects had a low SES, 48.8% a middle one, and 20.9% had a high SES. Our results revealed that a vegetarian diet is related to a lower BMI and less frequent alcohol consumption. Moreover, our results showed that a vegetarian diet is associated with poorer health (higher incidences of cancer, allergies, and mental health disorders), a higher need for health care, and poorer quality of life. Therefore, public health programs are needed in order to reduce the health risk due to nutritional factors.

Citation: Burkert NT, Muckenhuber J, Großschädl F, Rásky É, Freidl W (2014) Nutrition and Health – The Association between Eating Behavior and Various Health Parameters: A Matched Sample Study. PLoS ONE 9(2): e88278. <https://doi.org/10.1371/journal.pone.0088278>

Editor: Olga Y. Gorlova, Geisel School of Medicine at Dartmouth College, United States of America

Received: May 17, 2013; **Accepted:** January 9, 2014; **Published:** February 7, 2014

Copyright: © 2014 Burkert et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors have no support or funding to report.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Our diet has an impact on our well-being and on our health. Studies have shown a vegetarian diet to be associated with a lower incidence of hypertension, cholesterol problems, some chronic degenerative diseases, coronary artery disease, type II diabetes, gallstones, stroke, and certain cancers [1]–[7]. A vegetarian diet is characterized by a low consumption of saturated fat and cholesterol, due to a higher intake of fruits, vegetables and whole-grain products [3], [4], [8]. Overall, vegetarians have a lower body mass index [1], [4], [5], [7], [9]–[12], a higher socioeconomic status [13], and better health behavior, i.e. they are more physically active, drink less alcohol, and smoke less [9], [13], [14]. On the other hand, the mental health effects of a vegetarian diet or a Mediterranean diet rich in fruits, vegetables, whole-grain products and fish are divergent [9], [15]. For example, Michalak et al. [16] report that a vegetarian diet is associated with an elevated prevalence of mental disorders. A poor meat intake has been shown to be associated with lower mortality rates and higher life expectancy [17], and a diet which allows small amounts of red meat, fish and dairy products seems to be associated with a reduced risk of coronary heart disease as well as type 2 diabetes [18]. Additionally, evidence concerning lower rates of cancer, colon diseases including colon cancer, abdominal complaints, and all-cause mortality is, however, inconsistent [5]–[7], [19]–[22].

Not only is the intake of certain nutrients, like red meat, associated with an increased health risk [18], [23]–[26], high-caloric intake also plays a crucial role [23], [27]. Moreover, there seems to be proof that lifestyle factors like physical activity may be more crucial in lowering disease rates than individual dietary habits [20], [28]–[29]. While, generally speaking, diets based on plants, like vegetarian diets, seem to be associated with a certain health benefit, a lower risk to contract certain chronic diseases [30], and the ability to improve health [31]–[32], restrictive and monotonous vegetarian diets include the risk of nutritional deficits [2], [18], [19], [30], [33]. Baines et al. [9] report that vegetarians take more medication than non-vegetarians.

To summarize, a number of studies have shown vegetarian diets and diets with poor meat intake to be associated with lower mortality rates for certain diseases. Research about the dietary habits in Austria is, however, rather sparse and mainly focused on genetic factors [33]–[36]. Therefore, the aim of this study was to investigate health differences between different dietary habit groups among Austrian adults.

Methods

Study Design and Study Population

The sample for this cross-sectional study was taken from the Austrian Health Interview Survey (AT-HIS) which ran from March 2006 to February 2007 [37]. The AT-HIS is a standardized survey which is conducted at regular intervals in Austria (currently every eight years). The subjects included in the survey form a representative sample of the Austrian population. They were chosen from the central population register and are distributed across the different geographic regions of Austria. The AT-HIS is part of the European Health Interview Survey (E-HIS; <http://www.euhsid.org>), an important high-quality survey. The interviews were conducted by free-lancers engaged by the Austrian Statistic Agency. To ensure that all interviews were conducted in the same way, interviewers had to participate in a training day where they were instructed on how to conduct the survey. Time measurement, non-response analyses, and analyses of error dialogs were performed in order to ensure consistency between interviewers. Additionally, all interviewers were supervised by field supervisors. Overall, 15474 individuals, aged 15 years and older, were questioned in computer-assisted personal interviews (CAPI; 54.7% female; response rate: 63.1%).

While 0.2% of the interviewees were pure vegetarians (57.7% female), 0.8% reported to be vegetarians consuming milk and eggs (77.3% female), and 1.2% to be vegetarians consuming fish and/or eggs and milk (76.7% female). 23.6% reported to combine a carnivorous diet with lots of fruits and vegetables (67.2% female), 48.5% to eat a carnivorous diet less rich in meat (60.8% female), and 25.7% a carnivorous diet rich in meat (30.1% female). Since the three vegetarian diet groups included a rather small number of persons (N=343), they were analyzed as one dietary habit group. Moreover, since the vegetarian group was the smallest, we decided to match each of the vegetarians (1) with an individual of each other dietary habit group (carnivorous diet rich in fruits and vegetables (2), carnivorous diet less rich in meat (3) and a carnivorous rich in meat (4)).

Matching Process

In a first step, subjects consuming a vegetarian diet were identified (N=343). All vegetarians were categorized according to their sex, age (in age-groups spanning 5 years, e.g. 20- to 24-year-olds), and socioeconomic status (SES). Each such vegetarian was then matched with one subject consuming a carnivorous diet rich in fruits and vegetables, one individual eating a carnivorous diet less rich in meat, and one subject consuming a carnivorous diet rich in meat. Only 96.2% of the vegetarians were included in the analyses, since not all of them corresponded to a subject of the same sex, age, and SES from a different dietary habit group. Therefore, the total number of analyzed subjects was 1320 (comprising 330 vegetarians, 330 subjects consuming a carnivorous diet rich in fruits and vegetables, 330 individuals eating a carnivorous diet less rich in meat, and 330 subjects consuming a carnivorous diet rich in meat). Each dietary habit group was set-up according to the demographic characteristics shown in Table 1.

Age	SES	Sex		TOTAL	
		men	women	N	%
15-19	Low	3	22	25	7.6%
	Middle	9	19	28	8.5%
	High	0	1	1	0.3%
20-29	Low	4	11	15	4.5%
	Middle	10	35	45	13.6%
	High	5	13	18	5.4%
30-39	Low	3	10	13	3.9%
	Middle	6	23	29	8.8%
	High	6	11	17	5.2%
40-49	Low	4	6	10	3.0%
	Middle	4	20	24	7.3%
	High	8	16	24	7.3%
50-59	Low	3	7	10	3.0%
	Middle	3	11	14	4.2%
	High	1	6	7	2.1%
60-69	Low	3	7	10	3.0%
	Middle	1	14	15	4.5%
	High	0	2	2	0.6%
70-79	Low	4	8	12	3.6%
	Middle	1	2	3	0.9%
	High	0	0	0	0.0%
80+	Low	0	5	5	1.5%
	Middle	0	3	3	0.9%
	High	0	0	0	0.0%
TOTAL	Low	24	76	100	30.3%
	Middle	34	127	161	48.8%
	High	20	49	69	20.9%
TOTAL		78	252	330	100.0%

Note: Data source: Austrian Health Interview Survey (AT-HIS) 2006/07. N= number of subjects. Analyses were calculated with subjects matched according to their age, sex, and socio-economic status. Each dietary habit group consisted of 330 subjects with the above shown demographic characteristics (330 vegetarians, 330 subjects consuming a carnivorous diet rich in fruits and vegetables, 330 individuals eating a carnivorous diet less rich in meat, and 330 subjects consuming a carnivorous diet rich in meat; N= 1320). SES: A low SES corresponds a SES score between ≥ 3 and ≤ 6 , a middle SES between > 6 and ≤ 10 , and a high SES a score between > 10 and ≤ 15 .
doi:10.1371/journal.pone.0088278.t001

Table 1. Data definition and structure for each of the four dietary habit groups.

<https://doi.org/10.1371/journal.pone.0088278.t001>

Ethical Approval

The study was carried out in compliance with the principles laid down in the Helsinki Declaration. No minors or children were included in the study. Verbal informed consent was obtained from all subjects, witnessed, and formally recorded. The Ethics Committee of the Medical University of Graz approved the consent procedure as well as the conductance of this study (EK-number: 24-288 ex11/12).

Variables and Measurements

Face-to-face interviews were conducted by questioning the subjects about their socio-demographic characteristics, health-related behavior, diseases, medical treatments, and also psychological aspects.

The independent variable in this study was the dietary habit of individuals. Concerning eating behavior, the respondents were given a list of six different dietary habits and asked which one describes their eating behavior best (1=vegan, 2=vegetarian eating milk/eggs, 3=vegetarian eating fish and/or milk/eggs, 4=carnivorous diet rich in fruits and vegetables, 5=carnivorous diet less rich in meat, 6=carnivorous diet rich in meat). Participants described their dietary habit, without interviewers giving a clear definition of the various eating categories. Since, overall, only 2.2% of all participants consumed a vegetarian diet, these individuals were analyzed as one dietary habit group. We created a scale that would reflect the animal fat intake for each dietary habit (1=vegetarian diet, 2=carnivorous diet rich in fruits and vegetables, 3=carnivorous diet less rich in meat, 4=carnivorous diet rich in meat).

Since age, sex, and the socioeconomic background of subjects all have an influence on health [38]–[41], we matched the subjects according to these variables in order to control for their influence. The SES of the subjects (ranging between 3 and 15) was calculated using the following variables: net equivalent income, level of education, and occupation. Net equivalent income was calculated based on an equivalence scale provided by the OECD [42], and divided by quintiles. Level of education was measured by an ordinal variable, distinguishing between (1) basic education (up to 15 years of age), (2) apprenticeship/vocational school, (3) secondary education without diploma, (4) secondary education with diploma, and (5) university education. The occupation of the subjects was differentiated into the following five levels: (1) unskilled worker, (2) apprentice/skilled worker, (3) self-employed/middle job, (4) qualified job/academic, (5) executive position. To verify the combination of variables that served to calculate the SES, correlations with the different variables were calculated. They ranged between $r = .70$ and $r = .80$.

The body mass index (BMI) and lifestyle factors (physical activity, smoking, and alcohol consumption) were included as covariates in all analyses. The BMI was calculated by dividing the weight of a person in kilograms by the square of their height in meters (kg/m^2) [43]. Physical exercise was measured using the short version of the International Physical Activity Questionnaire (IPAQ), a self-reported instrument, which asks for an estimate of the total weekly physical activity (walking, moderate- and vigorous-intensity activity) performed during the last week. The short version of the IPAQ does not discriminate between leisure-time and non-leisure time physical activity. The total MET score was calculated by weighting the reported minutes per week within each activity by a MET energy expenditure estimate that was assigned to each category [44]. Smoking behavior was measured as the number of cigarettes smoked per day. Alcohol consumption was surveyed as the number of days on which alcohol was consumed during the last 28 days.

The dependent variables focusing on ill-health included self-perceived health, ranging from 1 (very good) to 5 (very bad), and impairment to health, ranging from 1 (very impaired) to 3 (not impaired). We further assessed the presence of 18 specific chronic conditions (asthma, allergies, diabetes, cataract, tinnitus, hypertension, cardiac infarction, apoplectic stroke, bronchitis, arthritis, sacrospinal complaints, osteoporosis, urinary incontinence, gastric or intestinal ulcer, cancer, migraine, mental illness (anxiety disorder or depression), and any other chronic condition). Each condition was coded as present (1) or absent (0). We calculated a total frequency score by summing up the chronic conditions present (0–18, sum index). Additionally, a vascular risk score was calculated by summing up the variables “hypertension”, “enhanced blood cholesterol level”, “diabetes”, and “smoking” (0–4, sum index). Each variable was coded as present (1) or absent (0).

A dependent variable concerning health care was created as the sum index of the number of doctors consulted in the last 12 months (0–8, sum index). Each of the 8 medical treatments (general practitioner, gynecologist, urologist, dermatologist, ophthalmologist, internist, orthopedist, and ENT physician) was coded as “consulted” (1) or “not consulted” (0). The number of vaccinations was analyzed by calculating a sum index combining 8 different vaccinations (influenza, tetanus, diphtheria, polio, FSME, pneumococci, hepatitis A and B; 0–8, sum index). Each vaccination was coded as present (1) or absent (0). In addition, preventive health care was analyzed by calculating a sum index of the variables “preventive check-ups”, “mammography”, “prostate gland check-up”, and “Papanicolaou test” (0–4, sum index). Each variable was coded as present (1) or absent (0).

The dependent variable concerning quality of life was measured using the short version of the WHOQOL (WHOQOL-BREF) [45]. Four domain scores (physical health, psychological health, social relationships, and environment) were calculated. These domain scores ranged between 4 and 20.

Statistical Analysis

In a first step subjects with different dietary habits (vegetarian, carnivorous diet rich in fruits and vegetables, carnivorous diet less rich in meat, carnivorous diet rich in meat) were matched according to their sex, age, and SES. Differences in lifestyle factors (BMI, total MET score, number of cigarettes smoked per day, and alcohol consumption in the last four weeks) between the different dietary habit groups were calculated by multivariate analysis of variance.

In order to analyze the differences between the dietary habit groups, multivariate analyses of variance were calculated for the three domains: (1) health (self-reported health, impairment due to health problems, number of chronic conditions, vascular risk), (2) health care (number of visits to the doctor, number of vaccinations, number of used preventive care offers), and (3) quality of life (physical and psychological health, social relationships, and environment). To address the bias of lifestyle factors impacting health, analyses of variance were calculated, controlling for the aforementioned lifestyle variables (BMI, physical activity, smoking behavior, and alcohol consumption).

In the domain of “health”, the two variables “self-reported health” and “impairment due to health problems” were originally assessed using an ordinal scale. Therefore, we controlled the results using non-parametric tests (Kruskal Wallis Test). Since the results were the same, only results of the analyses of variance are reported.

In addition, Chi-square tests were calculated for the aforementioned 18 chronic conditions in order to establish which one occurs significantly more often, depending on the form of nutrition. p -values $< .050$ were considered as statistically significant. All analyses were calculated using IBM SPSS software (version 20.0) for Windows.

Results

Participant Characteristics and Lifestyle Differences between the Dietary Habit Groups

In total, we analyzed the data of 1320 individuals (330 in each dietary habit group). Each dietary habit group was set-up according to the demographic characteristics shown in Table 1. Overall, 23.6% of all subjects were male and 76.4% female. 40.0% of the individuals were younger than 30 years, 17.8% between 30 and 39 years, 17.6% between 40 and 49 years, 9.4% between 50 and

59 years, 8.4% between 60 and 69 years, 4.4% between 70 and 79 years, and 2.4% than 80 years or older. 30.3% of the subjects had a low SES (they had an SES score of ≤ 6), 48.8% a middle one (SES between >6 and ≤ 10), and 20.9% had a high SES (SES >10).

Our multivariate analysis regarding lifestyle showed a significant main effect for the dietary habit of individuals ($p = .000$), showing that the different dietary habit groups differ in their overall health behavior. However, results of the univariate analyses showed that the dietary habit groups only differ concerning their BMI and their alcohol consumption.

Concerning BMI: vegetarians have the lowest mean BMI ($M = 22.9$), followed by subjects eating a carnivorous diet less rich in meat ($M = 23.4$), rich in fruits and vegetables ($M = 23.5$), and rich in meat ($M = 24.9$). Heavy meat eaters differ significantly from all other groups in terms of their BMI ($p = .000$).

Concerning physical exercise: no significant difference was found in the total MET score between the various dietary habit groups ($p = .631$).

Concerning smoking behavior: the number of cigarettes smoked per day did not differ between the various dietary habit groups ($p = .302$).

Concerning alcohol consumption: Subjects following a vegetarian diet ($M = 2.6$ days in the last 28 days) or a carnivorous diet rich in fruits and vegetables ($M = 3.0$ days) consume alcohol significantly less frequently than those eating a carnivorous diet less rich in meat ($M = 4.4$ days) or rich in meat ($M = 4.8$ days; $p = .000$).

Health Differences between the Dietary Habit Groups

In the domain of health, the multivariate analysis of variance showed a significant main effect for the dietary habit of individuals ($p = .000$). Overall, vegetarians are in a poorer state of health compared to the other dietary habit groups. Concerning self-reported health, vegetarians differ significantly from each of the other groups, toward poorer health ($p = .000$). Moreover, these subjects report higher levels of impairment from disorders ($p = .002$). Vegetarians additionally report more chronic diseases than those eating a carnivorous diet less rich in meat ($p = .000$; Table 2). Significantly more vegetarians suffer from allergies, cancer, and mental health ailments (anxiety, or depression) than the other dietary habit groups (Table 3). Subjects who eat a carnivorous diet rich in meat more often report urinary incontinence ($p = .023$). No differences between individuals consuming different forms of diet were found regarding their vascular risk ($p = .150$; Table 2).

Measure	Vegetarians (n = 100)		Carnivorous diet rich in fruits and vegetables (n = 100)		Carnivorous diet less rich in meat (n = 100)		Carnivorous diet rich in meat (n = 100)		p-value
	M	SD	M	SD	M	SD	M	SD	
Overall health (general)	1.76	0.94	1.73	0.93	1.74	0.91	1.67	0.91	<.001*
Impairment from disorders	1.69	0.99	1.71	1.03	1.73	0.99	1.73	1.01	<.001*
Chronic diseases	1.60	0.99	1.68	1.00	1.73	1.00	1.73	1.01	<.001*
Number of visits to the general practitioner	1.00	0.73	1.00	0.73	1.00	0.73	1.00	0.73	<.001*
Number of visits to the specialist	1.00	0.73	1.00	0.73	1.00	0.73	1.00	0.73	<.001*
Number of visits to the hospital	1.00	0.73	1.00	0.73	1.00	0.73	1.00	0.73	<.001*
Number of hospital days	1.00	0.73	1.00	0.73	1.00	0.73	1.00	0.73	<.001*
Number of hospital days (inpatient)	1.00	0.73	1.00	0.73	1.00	0.73	1.00	0.73	<.001*

Table 2. Differences in health and health care between the different dietary habit groups.
<https://doi.org/10.1371/journal.pone.0088278.t002>

Chronic condition	Vegetarian	Carnivorous diet rich in fruits and vegetables	Carnivorous diet less rich in meat	Carnivorous diet rich in meat	p-value (χ^2)
Allergies	14.0%	7.0%	3.0%	4.0%	.272
Arteriosclerosis	18.0%	18.0%	20.0%	18.0%	.600
Diabetes	2.0%	4.0%	2.0%	2.0%	.451
Cancer	4.0%	3.0%	3.0%	3.0%	.340
Stroke	4.0%	4.0%	4.0%	3.0%	.261
Hyperuricemia	11.0%	10.0%	12.0%	11.0%	.261
Urinary incontinence	1.0%	1.0%	2.0%	1.0%	.202
Psychiatric disorder	1.0%	1.0%	1.0%	1.0%	.410
Benign prostatic hyperplasia	3.0%	3.0%	2.0%	3.0%	.701
Arthritis	6.0%	7.0%	4.0%	10.0%	.022
Cardiovascular complications	24.0%	24.0%	18.0%	20.0%	.060
Hypertension	6.0%	4.0%	3.0%	3.0%	.411
Urinary incontinence	2.0%	3.0%	2.0%	4.0%	.202
Stroke or myocardial infarction	4.0%	4.0%	1.0%	3.0%	.101
Cholesterol	14.0%	13.0%	13.0%	14.0%	.202
Insomnia	10.0%	11.0%	9.0%	11.0%	.074
Mental stress (anxiety, depression)	8.0%	4.0%	5.0%	4.0%	.034
Any other chronic conditions	8.0%	5.0%	5.0%	6.0%	.301

Table 3. Differences in suffering from various chronic conditions between the different dietary habit groups.
<https://doi.org/10.1371/journal.pone.0088278.t003>

Differences in Health Care between the Dietary Habit Groups

Our multivariate analysis regarding health care has shown a significant main effect for dietary habits ($p = .000$) and confirmed that, overall, subjects with a lower animal fat intake demonstrate worse health care practices. Vegetarians and subjects eating a carnivorous diet rich in fruits and vegetables consult doctors more often than those eating a carnivorous diet less rich in meat ($p = .003$). Moreover, vegetarians are vaccinated less often than all other dietary habit groups ($p = .005$) and make use of preventive check-ups less frequently than subjects eating a carnivorous diet rich in fruits and vegetables ($p = .033$; Table 2).

Differences in Quality of Life between the Dietary Habit Groups

Regarding quality of life, the main effect of the multivariate analysis of variance showed no significant difference between the dietary habit groups ($p = .291$). The results obtained in the univariate analyses of variance, however, revealed that vegetarians have a lower quality of life in the domains of "physical health" ($p = .026$) and "environment" ($p = .037$) than subjects consuming a carnivorous diet less rich in meat. Moreover, vegetarians have a lower quality of life regarding "social relationships" than individuals eating a carnivorous diet rich in fruits and vegetables, or those with a carnivorous diet less rich in meat ($p = .043$). All results are shown in Table 4.

Variable	Vegetarian (n=122)		Semi-vegetarian (n=122)		Semi-carnivorous (n=122)		Carnivorous (n=122)		p-value
	n	Mean	n	Mean	n	Mean	n	Mean	
Number of visits	122	1.0	122	1.0	122	1.0	122	1.0	NS
Number of visits (range)	122	1-10	122	1-10	122	1-10	122	1-10	NS
Number of visits (SD)	122	1.0	122	1.0	122	1.0	122	1.0	NS
Number of visits (IQR)	122	1.0	122	1.0	122	1.0	122	1.0	NS
Number of visits (95% CI)	122	0.9-1.1	122	0.9-1.1	122	0.9-1.1	122	0.9-1.1	NS

Table 4. Differences in quality of life between the different dietary habit groups.
<https://doi.org/10.1371/journal.pone.0088278.t004>

Discussion

Overall, our findings reveal that vegetarians report poorer health, follow medical treatment more frequently, have worse preventive health care practices, and have a lower quality of life. Concerning the variable “eating behavior”, we tried to generate a variable that would reflect the animal fat intake (1=vegetarian, 2=carnivorous diet rich in fruits and vegetables, 3=carnivorous diet less rich in meat, 4=carnivorous diet rich in meat). The mean BMI of subjects is coupled in nearly linear progression with the amount of animal fat intake. This is in line with previous studies showing vegetarians to have a lower body mass index [1], [4], [5], [7], [9]–[12].

Our results have shown that vegetarians report chronic conditions and poorer subjective health more frequently. This might indicate that the vegetarians in our study consume this form of diet as a consequence of their disorders, since a vegetarian diet is often recommended as a method to manage weight [10] and health [46]. Unfortunately, food intake was not measured in more detail, e.g. caloric intake was not covered. Hence, further studies will be necessary to analyze health and its relationship with different forms of dietary habits in more detail.

When analyzing the frequency of chronic diseases, we found significantly higher cancer incidence rates in vegetarians than in subjects with other dietary habits. This is in line with previous findings, reporting that evidence about cancer rates, abdominal complaints, and all-cause mortality in vegetarians is rather inconsistent [5]–[7], [19]–[22]. The higher cancer incidence in vegetarians in our study might be a coincidence, and is possibly related to factors other than the general amount of animal fat intake, such as health-conscious behavior, since no differences were found regarding smoking behavior and physical activity in Austrian adults as reported in other studies for other countries [9], [13], [14]. Therefore, further studies will be required in Austria in order to analyze the incidence of different types of cancer and their association with nutritional factors in more depth.

Several studies have shown the mental health effects of a vegetarian diet to be divergent [9], [15], [16]. Vegetarians in our study suffer significantly more often from anxiety disorder and/or depression. Additionally, they have a poorer quality of life in terms of physical health, social relationships, and environmental factors.

Moreover, the use of health care differs significantly between the dietary habit groups in our study. Vegetarians need more medical treatment than subjects following another form of diet. However, this might be due to the number of chronic conditions, which is higher in subjects with a vegetarian diet.

Among the **strengths** of our study are: the large sample size, the matching according to age, sex, and socioeconomic background, and the standardized measurement of all variables. Other strengths of our study include considering the influence of weight and lifestyle factors on health, e.g. physical exercise and smoking behavior.

Potential **limitations** of our results are due to the fact that the survey was based on cross-sectional data. Therefore, no statements can be made whether the poorer health in vegetarians in our study is caused by their dietary habit or if they consume this form of diet due to their poorer health status. We cannot state whether a causal relationship exists, but describe ascertained associations. Moreover, we cannot give any information regarding the long-term consequences of consuming a special diet nor concerning mortality rates. Thus, further longitudinal studies will be required to substantiate our results. Further limitations include the measurement of dietary habits as a self-reported variable and the fact that subjects were asked how they would describe their eating behavior, without giving them a clear definition of the various dietary habit groups. However, a significant association between the dietary habit of individuals and their weight and drinking behavior is indicative for the validity of the variable. Another limitation concerns the lack of detailed information regarding nutritional components (e.g. the amount of carbohydrates, cholesterol, or fatty acids consumed). Therefore, more in-depth studies about nutritional habits and their effects on health are required among Austrian adults. Further studies should e.g. investigate the influence of the various dietary habits on the incidence of different cancer types. To our knowledge this is the first study ever in Austria to analyze differences in terms of dietary habits and their impact on health. We admit that the large number of participants made it necessary to keep the questions simple, in order to cover the large sample. Overall, we feel that our results are of specific interest and contribute to extant scientific knowledge, notwithstanding some limitations regarding causes and effects.

Conclusions

Our study has shown that Austrian adults who consume a vegetarian diet are less healthy (in terms of cancer, allergies, and mental health disorders), have a lower quality of life, and also require more medical treatment. Therefore, a continued strong public health program for Austria is required in order to reduce the health risk due to nutritional factors. Moreover, our results emphasize the necessity of further studies in Austria, for a more in-depth analysis of the health effects of different dietary habits.

Author Contributions

Conceived and designed the experiments: NTB WF. Performed the experiments: NTB. Analyzed the data: NTB JM FG ER WF. Contributed reagents/materials/analysis tools: NTB JM FG WF. Wrote the paper: NTB ER WF.

References

1. American Dietetic Association, Dietitians of Canada (2003) Position of the American Dietetic Association and Dietitians of Canada: vegetarian diets. *Can J Diet Pract Res* 64(2): 62–81.
View Article • PubMed/NCBI • Google Scholar
2. Craig WJ (2009) Health effects of vegan diets. *Am J Clin Nutr* 89(5): 1627s–1633s.
View Article • PubMed/NCBI • Google Scholar

3. Craig WJ (2010) Nutrition concerns and health effects of vegetarian diets. *Nutr Clin Pract* 25(6): 613–20.
View Article • PubMed/NCBI • Google Scholar
4. Craig WJ, Mangels AR (2009) American Dietetic Association (2009) Position of the American Dietetic Association: vegetarian diets. *J Am Diet Assoc* 109(7): 1266–82.
View Article • PubMed/NCBI • Google Scholar
5. Dwyer JT (1988) Health aspects of vegetarian diets. *Am J Clin Nutr* 48: 712–38.
View Article • PubMed/NCBI • Google Scholar
6. Key TJ, Appleby PN, Davey GK, Allen NE, Spencer EA, et al. (2003) Mortality in British vegetarians: review and preliminary results from EPIC-Oxford. *Am J Clin Nutr* 78: 533s–58s.
View Article • PubMed/NCBI • Google Scholar
7. Key TJ, Appleby PN, Rosell MS (2006) Health effects of vegetarian and vegan diets. *Proc Nutr Soc* 65(1): 35–41.
View Article • PubMed/NCBI • Google Scholar
8. Deriemaeker P, Alewaeters K, Hebbelinc M, Lefvre J, Philippaerts R, et al. (2010) Nutritional status of Flemish vegetarians compared with non-vegetarians: a matched sample study. *Nutrients* 2(7): 770–80.
View Article • PubMed/NCBI • Google Scholar
9. Baines S, Powers J, Brown WJ (2007) How does the health and well-being of young Australian vegetarian women compare with non-vegetarians? *Public Health Nutr* 10(5): 436–42.
View Article • PubMed/NCBI • Google Scholar
10. Farmer B, Larson BT, Fulgoni VL, Rainville AJ, Liepa GU (2011) A vegetarian dietary pattern as a nutrient-dense approach to weight management: an analysis of the national health and nutrition examination survey 1999–2004. *J Am Diet Assoc* 111(6): 819–27.
View Article • PubMed/NCBI • Google Scholar
11. Rizzo NS, Sabaté J, Jaceldo-Siegl K, Fraser GE (2011) Vegetarian dietary patterns are associated with a lower risk of metabolic syndrome: the adventist health study 2. *Diabetes Care* 34(5): 1225–27.
View Article • PubMed/NCBI • Google Scholar
12. Tonstad S, Butler T, Yan R, Fraser GE (2009) Type of vegetarian diet, body weight, and prevalence of type 2 diabetes. *Diabetes care* 32(5): 791–96.
View Article • PubMed/NCBI • Google Scholar
13. Pollard J, Greenwood D, Kirk S, Cade J (2001) Lifestyle factors affecting fruit and vegetable consumption in the UK women's cohort study. *Appetite* 37(1): 71–79.
View Article • PubMed/NCBI • Google Scholar
14. Gacek M (2010) Selected lifestyle and health condition indices of adults with varied models of eating. *Rocz Pantstw Zakl Hig* 61(1): 65–69.
View Article • PubMed/NCBI • Google Scholar
15. Low Dog T (2010) The role of nutrition in mental health. *Altern Ther Health Med* 16(2): 42–46.
View Article • PubMed/NCBI • Google Scholar
16. Michalak J, Zhang XC, Jacobi F (2012) Vegetarian diet and mental disorders: results from a representative community survey. *Int J Behav Nutr Phys Act* 9: 67.
View Article • PubMed/NCBI • Google Scholar
17. Singh PN, Sabaté J, Fraser GE (2003) Does low meat intake increase life expectancy in humans? *Am J Clin Nutr* 78(3): 526s–32s.
View Article • PubMed/NCBI • Google Scholar
18. McEvoy CT, Temple N, Woodside JV (2012) Vegetarian diets, low-meat diets and health: a review. *Public Health Nutr* 15: 2287–2294.
View Article • PubMed/NCBI • Google Scholar
19. Appelby PN, Thorogood M, Mann JI, Key TJA (1999) The Oxford vegetarian study: an overview. *Am J Clin Nutr* 70: 525s–31s.
View Article • PubMed/NCBI • Google Scholar
20. Frentzel-Beyme R, Chang-Claude J (1994) Vegetarian diets and colon cancer: the German experience. *Am J Clin Nutr* 59: 143s–52s.
View Article • PubMed/NCBI • Google Scholar
21. Burkert NT, Großschädl F, Muckenhuber J, Rásky É, Stronegger WJ, et al. (2012) Gastro-intestinal complaints related to various eating behaviours. *J Nutr Therapeutics* 1: 19–23.
View Article • PubMed/NCBI • Google Scholar
22. Tantamango-Bartley Y, Jaceldo-Siegl K, Fan J, Fraser G (2013) Vegetarian Diets and the Incidence of Cancer in a Low-risk Population. *Cancer Epidemiol Biomarkers Prev* 22(2). DOI:10.1158/1055-9965.EPI-12.1060.

23. Flood A, Velie AM, Sinha R, Chatterjee N, Lacey JV, et al. (2003) Meat, fat, and their subtypes as risk factors for colorectal cancer in a prospective cohort study in women. *Am J Epidemiol* 158: 59–68.
View Article • PubMed/NCBI • Google Scholar
24. Mathew A, Peters U, Chatterjee N, Kulldorff M, Sinha R (2004) Fat, fiber, fruits, vegetables, and risk of colorectal adenomas. *Int J Cancer* 108: 287–292.
View Article • PubMed/NCBI • Google Scholar
25. Seow A, Quah SR, Nyam D, Straughan PT, Chua T, et al. (2002) Food groups and the risk of colorectal carcinomas in an Asian population. *Cancer* 95: 2390–2396.
View Article • PubMed/NCBI • Google Scholar
26. Vang A, Singh PN, Lee JW, Haddad EH, Brinegar CH (2008) Meats, processed meats, obesity, weight gain and occurrence of diabetes among adults: findings from Adventist Health Studies. *Ann Nutr Metab* 52(2): 96–104.
View Article • PubMed/NCBI • Google Scholar
27. Boutron-Ruault MC, Senesse P, Meance S, Belghiti C, Faivre J (2001) Energy intake, body mass index, physical activity, and the colorectal adenoma-carcinoma sequence. *Nutr Cancer* 39: 50–57.
View Article • PubMed/NCBI • Google Scholar
28. Appleby PN, Key TJ, Thorogood M, Burr ML, Mann J (2002) Mortality in British vegetarians. *Public Health Nutr* 5(1): 29–36.
View Article • PubMed/NCBI • Google Scholar
29. Chang-Claude J, Hermann S, Eilber U, Steindorf K (2005) Lifestyle determinants and mortality in German vegetarians and health-conscious persons: results of a 21-year follow-up. *Cancer Epidemiol Biomarkers Prev* 14(4): 963–68.
View Article • PubMed/NCBI • Google Scholar
30. Fraser GE (2009) Vegetarian diets: what do we know of their effects on common chronic diseases? *Am J Clin Nutr* 89: 1607s–12s.
View Article • PubMed/NCBI • Google Scholar
31. Sabaté J (2003a) The contribution of vegetarian diets to health and disease: a paradigm shift? *Am J Clin Nutr* 78: 502s–7s.
View Article • PubMed/NCBI • Google Scholar
32. Sabaté J (2003b) The contribution of vegetarian diets to human health. *Forum Nutr* 56: 218–20.
View Article • PubMed/NCBI • Google Scholar
33. Karlic H, Schuster D, Varga F, Klindert G, Lapin A, et al. (2008) Vegetarian diet affects genes of oxidative metabolism and collagen synthesis. *Ann Nutr Metab* 53(1): 29–32.
View Article • PubMed/NCBI • Google Scholar
34. Kornsteiner M, Singer I, Elmadfa I (2008) Very low n-3 long-chain polyunsaturated fatty acid status in Austrian vegetarians and vegans. *Ann Nutr Metab* 52(1): 37–47.
View Article • PubMed/NCBI • Google Scholar
35. Majchrzak D, Singer I, Maenner M, Rust P, Genser D, et al. (2006) B-vitamin status and concentrations of homocysteine in Austrian omnivores, vegetarians and vegans. *Ann Nutr Metab* 50(6): 485–91.
View Article • PubMed/NCBI • Google Scholar
36. Schumacher M, Eber B, Schallmoser K, Toplak H, Zweiker R, et al. (1993) Eating behavior of patients with metabolic diseases and metabolically healthy probands in Austria. Results of a questionnaire survey at the Graz Autumn Fair 1991. *Wien Med Wochenschr* 43(12): 329–23.
View Article • PubMed/NCBI • Google Scholar
37. Klimont J, Kytir J, Leitner B (2007) Austrian Health Interview Survey 2006/07. (Oesterreichische Gesundheitsbefragung 2006/07. Hauptergebnisse und methodische Dokumentation.) Wien: Statistik Austria.
38. Bauer GF, Huber CA, Jenny GJ, Mueller F, Haemming O (2009) Socioeconomic status, working conditions and self-rated health in Switzerland: explaining the gradient in men and women. *International J Pub Health* 54: 23–30.
View Article • PubMed/NCBI • Google Scholar
39. Volkert D (2006) The body mass index (BMI) – an important parameter to evaluate the nutritional status [Der Body-Mass-Index (BMI) – ein wichtiger Parameter zur Beurteilung des Ernährungszustandes]. *Aktuel Ernahr Med* 31: 126–132.
View Article • PubMed/NCBI • Google Scholar
40. Dörner TE, Rieder A (2010) Obesity paradox or reverse epidemiology: is high body weight a protective factor for various chronic conditions? [Das Adipositasparadoxon oder Reverse Epidemiologie. Hohes Körpergewicht als protektiver Faktor bei bestimmten chronischen Bedingungen?] *Dtsch Med Wochenschr* 135: 413–418.
View Article • PubMed/NCBI • Google Scholar
41. Burkert NT, Rásky E, Freidl W (2012) Social inequalities regarding health and health behaviour in Austrian adults. *Wien klin Wochenschr* 7–8: 256–261.
View Article • PubMed/NCBI • Google Scholar
42. OECD Social Policy Division (2009) “What are equivalence scales?” Available: <http://www.oecd.org/els/social>.

43. World Health Organization (2013) Global database on Body Mass Index. BMI classification. Available: http://apps.who.int/bmi/index.jsp?introPage=intro_3.html. Accessed 2013 September 17.
44. Craig CL, Marshall A, Sjoström M, Bauman AE, Booth ML, et al. (2003) International Physical Activity Questionnaire: 12 country reliability and validity. *Med Sci Sports Exerc* 35(8): 1381–1395.
View Article • PubMed/NCBI • Google Scholar
45. World Health Organization (1996) WHOQOL-BREF. Introduction, Administration, Scoring, and Generic Version of the Assessment. Geneva: WHO.
Available: http://www.who.int/mental_health/media/en/76.pdf.
46. Leitzmann C (2005) Vegetarian diets: what are the advantages? *Forum Nutr* 57: 147–56.
View Article • PubMed/NCBI • Google Scholar